

Technical Supplement: Emergency Department Patient Survey, 2014-15

February 2016

Revision History

Version	Issue Date	Author	Comments
1.0	March 2016	Jason Boyd	Second EDPS technical supplement released – updated for the 2014-15 survey year

Contents

The NSW Patient Survey Program	1
The Emergency Department Patient Survey	2
Organisational roles in producing survey samples	3
Inclusion criteria	4
Phase 1 screening	4
Phase 2 screening	5
Drawing of the sample	5
Data Management	8
Data Analysis	9
Completeness of survey questionnaires	9
Calculation of response rate	9
Weighting of data	9
Demographic characteristics of respondents to EDPS	9
Reporting	15
Confidentiality	17
Statistical Analysis	17
Calculation of percentages	19
Appendix 1	20
Facilities included in the EDPS sampling frame	20
Appendix 2	22
Missing and 'Don't know' responses	22
Appendix 3	26
Dariyad magauraa	26

Suggested Citation: Technical Supplement: Emergency Department Patient Survey, 2014-15. Bureau of Health Information. Sydney (NSW); 2016.

Date of publication: March 2016

Please note that there is the potential for minor revisions of information in this report. Please check the online version at www.bhi.nsw.gov.au for any amendments.

The NSW Patient Survey Program

The NSW Patient Survey Program began surveying patients in NSW public facilities from 2007. From 2007 to mid-2012, the program was co-ordinated by the NSW Ministry of Health using questionnaires obtained under license from NRC Picker. Ipsos Social Research Institute Ltd (Ipsos) was contracted to manage the logistics of the survey program. Responsibility for the Patient Survey Program was transferred from the Ministry of Health to the Bureau of Health Information (BHI) in July 2012, with Ipsos continuing as the contracted partner to manage the logistics.

The aim of the program is to measure and report on patients' experiences of care in public health facilities in New South Wales (NSW), on behalf of the NSW Ministry of Health and the local health districts (LHDs). The results are used as a source of performance measurement for individual hospitals, LHDs and NSW as a whole.

This document outlines the sampling methodology, data management and analysis of the 2014-15 Emergency Department Patient Survey (EDPS).

For more information on how to interpret results and statistical analysis of differences between hospitals, LHDs or NSW, please refer to the "Guide to Interpreting Differences" at www.bhi.nsw.gov.au/nsw_patient_survey_program.

The Emergency Department Patient Survey

In 2013, the EDPS was the second survey to be sent to patients as part of the revised NSW Patient Survey Program, after the Adult Admitted Patient Survey. In 2014, the EDPS was conducted for the second time, covering the period of April 2014 to March 2015.

Significant changes were made to the questionnaire content between the 2013-14 and 2014-15 questionnaire versions. These changes were made to improve navigation through the questionnaire and in response to the latest stakeholder needs. These changes were informed by an analysis of 2013/14 results, specifically item non-response to survey questions, percentage of invalid responses to questions, floor and ceiling effects (based on the mean, standard deviation and skewness of results), and correlation to other questions in the questionnaire.

Changes were also made to the sampling for the survey due to lower response rates seen for younger patients in the EDPS. In order to ensure that sufficient numbers of younger patients were responding to allow reporting, the proportion of younger patients sampled in the survey was increased. It was expected that, while this would have a negative impact on unadjusted response rates, it would improve the overall representativeness of respondents to the survey.

Organisational roles in producing survey samples

The survey program assures patients that their responses will be confidential and that staff at hospitals will not be able to determine who gave which response. BHI does this through a number of mechanisms, including:

- Data suppression (results for fewer than 30 responses are suppressed)
- Reporting aggregated results
- De-identification of patient comments
- Segregation of roles when constructing the survey samples (see below).

The sampling method for the NSW Patient Survey Program requires collaboration between staff at BHI, Ipsos SRI and the Ministry of Health's Health Systems Performance Information and Reporting Branch (HSPIRB) (see Figure 1). This survey used data obtained from the Health Information Exchange (HIE).

BHI has access to confidentialised unit record data from selected tables of the HIE database. Use of an encrypted patient number allows deduplication at the patient level within a hospital. For the EDPS, sampling frames are defined separately for each month, with the date of emergency department (ED) attendance is used to define eligible records. Sample sizes for each included hospital are calculated in advance, as defined later in this report.

Figure 1: Organisational responsibilities in sampling and survey processing, Emergency Department Patient Survey, 2014-15

BHI

- Determine inclusion and exclusion rules in association with stakeholders
- Develop sampling strategy including strata and included facilities based on requests from stakeholders and availability of data in the database available for sampling (HIE in the case of admitted patient surveys)
- · Calculate target sample sizes by strata within facilities and provide to HSIPRB
- Extract monthly data from HIE, create interim sampling frame following phase 1 screening and send via secure file transfer to HSIPRB

HSIPRB

- · Add names and addresses to interim sampling frame
- Undergo phase 2 cleaning and exclusions
- Generate samples based on sampling targets provided by BHI
- Provide mailing list via secure file transfer to Ipsos

Ipsos

- · Administer the survey fieldwork, collate results, clean results
- Provide datafile of results to BHI for analysis, via secure file transfer, once all name and address information is removed

Inclusion criteria

Phase 1 screening

Emergency department patient data pass through two phases of cleaning. The first phase of screening is applied by BHI. Many of these criteria are developed in conjunction with advice of stakeholders.

Inclusions

• Patients who visited an emergency department in a NSW public hospital with a peer group classification of A1, A2, A3, B, C1 or C2.

Exclusions

 Patients who were dead on arrival or died in ED (mode of separation of 8 and 3 respectively) were excluded from the sample.

A series of further exclusion criteria were applied to take into account a range of factors including: the potentially high vulnerability of particular patient groups and/or patients with particularly sensitive reasons for admission; certain patients' ability to answer questions about their experiences; and the relevance of the survey questions to particular patient groups.

The effectiveness of this screening is reduced for the EDPS compared to the Adult Admitted Patient Survey (AAPS) due to the variables in the dataset. For example, the ED dataset does not contain robust diagnosis (ICD-10-AM) information that allows these exclusions. Because of this, further screening to exclude sensitive groups can only be done for patients subsequently admitted to hospital. In addition, ED patients subsequently admitted to hospital (mode of separation of 1,10,11,12 or 13) with the following procedures or diagnoses that were recorded for their inpatient stay were omitted:

- admitted for a termination of pregnancy procedure [35643-03];
- treated for maltreatment syndromes [T74] in any diagnosis field, including neglect or abandonment, physical abuse, sexual abuse, psychological abuse, other maltreatment syndromes and maltreatment syndrome, unspecified;
- treated for contraceptive management [Z30] in any diagnosis field, including general counselling and advice on contraception, surveillance of contraceptive drugs, surveillance of contraceptive device, other contraceptive management and contraceptive management, unspecified;
- patients with a diagnosis of stillborn baby [Z37] in any diagnosis field (including single stillbirth, twins, one liveborn and one stillborn, twins, both stillborn and other multiple births, some liveborn) were excluded.
- In addition, where ED patients were admitted to hospital, they were excluded if in the subsequent admission they had a mode of separation of death.

From October 2014, the following additional exclusions were applied:

- Intentional self-harm: ICD10 code between X60 and X84
- Sequelae of intentional self-harm: ICD10 code = Y87.0
- Unspecified event, undetermined intent: ICD10 code commences with Y34
- Suicidal ideation: ICD10 code = R45.81
- Family history of other mental and behavioural disorders: ICD10 code commences with Z81.8

Personal history of self-harm: ICD10 code commences with Z91.5.

Where patients had multiple visits within the sampling month, their most recent hospital visit was kept. The questionnaire asks patients to respond to the survey based on their most recent ED visit in a particular month.

Phase 2 screening

BHI provides the interim sampling frame to HSIPRB, who add patient name and address information. Data then undergo a second phase of screening. This involves exclusions for administrative/logistical reasons, or where death had been recorded after discharge for the stay used for sample selection but before the final sampling frame is prepared.

Exclusions

- Invalid address (including those with addresses listed as hotels, motels, nursing homes, Community Services, Mathew Talbot hostel, 100 William Street, army quarters, jails, unknown, NFA)
- Invalid name (including twin, baby of, etc.)
- Invalid date of birth
- On the 'do not contact' list
- Sampled in the previous six months for any BHI patient survey currently underway
- Had a death recorded according to the NSW Birth Deaths and Marriages Registry and/or Agency Performance and Data Collection, prior to the sample being provided to Ipsos.

The data following these exclusions are defined by BHI as the final sampling frame.

Drawing of the sample

Survey design

A stratified sample design was applied, with each facility defined as a stratum. Within each facility, patients are further stratified by the following variables:

• Age – aged 0-17, 18-49 or 50 years and over, based on the age variable.

Although sampling is undertaken monthly, sample size calculations are based on whether reporting is on a quarterly or annual basis. All facilities in C1 or C2 peer groups were sampled for annual reporting, whereas facilities in the remaining peer groups were sampled for quarterly reporting. For the purposes of sampling, the population of Sydney and Sydney Eye Hospitals were combined. In addition:

- all patients at the two children's hospitals were included in the 'under 18' stratum for sampling purposes
- children under 18 years admitted to A3 (Ungrouped Acute tertiary referral) facilities were included in the '18 to 49' age stratum because of very small numbers in the under 18 age group for these three hospitals.

Patients were selected within strata using simple random sampling without replacement. Sample sizes were defined at the facility level (or by patient type as described within the next section), with proportional sampling of strata within facilities/patient types.

The monthly targets by strata for the 2014-15 sampling period were based on the emergency department patient data from 2013 (after Phase 1 of the screening process).

The required sample size for each facility (i) within reporting stratum (j) was estimated using Equation 1.

Equation 1

$$s_{ij} = \frac{\chi^2 N_{ij} P(1 - P)}{d^2(N_{ij} - 1) + \chi^2 P(1 - P)}$$

Where:

 s_{ii} = estimated sample size for facility i and stratum j

 χ^2 = tabulated value of chi-squared with one degree of freedom at 5% level of significance (3.841)

 N_{ij} = population in the reporting stratum j of facility i, estimated using data from the 2013 calendar year with phase 1 exclusion criteria applied, aggregated to correspond with the reporting period (i.e. by quarter or full year)

P = expected proportion giving the most positive response to the question on satisfaction with overall care (0.8), based on previous levels of response to patient surveys

d = degree of accuracy of the 95% confidence interval expressed as a proportion (±0.07).

The sample size calculation aimed for a confidence interval around an expected proportion of 0.8 of ± 0.07 at the reporting strata level within each facility. Sample sizes were then allocated proportionately across strata internal to these reporting strata.

Finally, cell sample sizes are increased to account for fewer than 100% of patients responding to the survey. This was done by dividing the expected sample size by the expected response rate. Response rates for each stratum was estimated based on response rates observed in the 2013-14 survey (Table 1):

Table 1: Response rates used when calculating the targets for mailing, AAPS 2014

Stratum	Quarterly reporting (A and B peer groups)	Annual reporting (C peer groups)
0-17 years	25%	25%
18-49 years	25%	20%
50+ years	55%	50%

In addition, monthly mailing targets were changed from variable (depending on expected monthly patient numbers) to fixed monthly numbers (based on annual sample sizes divided by 12 months). A minimum monthly target of four patients was applied to all strata (e.g. if calculations require one, two or three patients in any stratum, this will be increased to four patients).

The adjusted sample sizes file was provided to HSIPRB as the targets for the 2014-15 survey. For each month of sampling, HSPIRB randomly selected patients within each stratum, according to mailing targets provided by BHI.

Notes:

- The sample size calculation based on Equation 1 (page 7) assumes simple random sampling, whereas a stratified survey design was used. This, and differences in the response rate between strata, may result in some estimates having wider confidence intervals than expected, even when the prevalence is 80%.
- For the purposes of sampling and reporting, the population of Sydney and Sydney Eye Hospitals were combined as one facility.

Data Management

Data collection

Upon completion of a survey questionnaire, the respondent returns or submits the completed survey (depending on whether they completed the paper-based questionnaire or the online questionnaire) to Ipsos. If a paper form is returned, Ipsos then scans in the answers electronically and manually enters free text fields.

Once all of the data is collated into a single dataset, all names and addresses are removed from the dataset. Also, all text entry fields are checked for potential identifiers (names of patients, names of doctors, telephone numbers, etc.) and any that are found are replaced with "XXXX".

Following this, each record is checked for any errors in completion and reasonable adjustments (known as 'cleaning') are made to the dataset, for example, removing responses where the patient has not correctly followed questionnaire instructions or providing multiple answers to a single response question.

At the end of this process, Ipsos uses a secure NSW Ministry of Health system to transfer the data from their servers to BHI's secure servers, all of which are password protected with limited staff access.

At no stage do BHI, who analyse the data, have access to the names and contact details of the respondents. This ensures respondent answers remain confidential and identifying data can never be publicly released.

Data Analysis

Completeness of survey questionnaires

In EDPS 2014-15, the completeness of responses was very high, with 99% of respondents answering more than 77 questions and 95% answering more than 79 questions, out of the 93 questions in the questionnaire. Fifteen respondents answered none of the questions, with the exception of the two free-text questions at the end of the questionnaire – these respondents were excluded from further analysis.

Calculation of weighted response rate

The response rate is the proportion of people sampled in the survey that actually completed and returned their survey form. As a result of the oversampling of younger patients, the distribution of patients in the sample (patients who were sent questionnaires) does not match the age distribution of patients in the population (Table 2). Therefore, response rates were adjusted to ensure that the overall survey response rate reflects a response rate that would be observed if patients were sampled proportional to the patient mix, creating the 'weighted response rate'. The weighted response rates are shown in Tables 4 and 5 in the following sections.

Table 2: Patient population distribution and corresponding number of surveys mailed, EDPS 2014-15

Age group	Percentage in patient pop	Percentage in sample	Percentage in respondents
0-17	26%	32%	32%
18-49	39%	50%	30%
50+	35%	19%	38%

Weighting of data

The protocol of the NSW Patient Survey Program is, when possible, to 'weight' data to account for differences (bias) in the probability of sampling and the likelihood of different patient groups to respond. Weighting makes the results more representative of the overall patient population, making the data more useful for the purposes of decision-making and service improvement.

Weights were calculated in two stages. Weights are calculated for each quarter of data as they become available. Once 12 months of data were available, weights for facilities reported on an annual basis was adjusted, to better reflect patient populations (which was difficult to do due to smaller numbers of respondents at the quarterly level).

Weighting of quarterly data

For each quarter of data, responses were weighted to match the population by age (Under 18, 18–49 or 50+ years) and visit type (admitted or non-admitted emergency).

Data were weighted at facility level for hospitals that were sampled for quarterly reporting (peer group hospitals A1, A2, A3 and B) and at LHD level for hospitals that were sampled for annual reporting (peer group hospitals C1 and C2). Methods for weighting are described in the following pages.

Calculating quarterly response weights

Interim quarterly response weights were calculated as:

$$w_{ij} = \frac{N_{ij}}{n_{ii}} \tag{1}$$

where:

 N_{ij} denotes the population (i.e. total number of patients eligible for the survey) of the i^{th} facility in the j^{th} age group. Eligible patient numbers were based on the number of patients following the second phase of screening undertaken by the Ministry of Health.

 n_{ij} denotes the sample size (i.e. number of respondents) of the i^{th} facility in the j^{th} age group.

If the stratum cell size within a facility was five or fewer, then cells within that facility were aggregated for weighting purposes by grouping across age group.

The interim quarterly weights were then passed through the GREGWT macro, a survey-specific SAS program developed by the ABS to assist with weighting of complex survey data¹. It uses iterative proportional fitting to ensure that the weights at the margins agreed with the population totals even though it is often impossible for the weights to equal the population at the individual cell level. The marginal totals specified were:

- Benchmark 1: Facility (with annually-reported facilities within the same LHD combined)
- Benchmark 2: Peer group
- Benchmark 3: Peer group (with C1 and C2 facilities combined) x age (with some strata combined see below)
- Benchmark 4: Peer group (with C1 and C2 facilities combined) x visit type

¹ Bell, P. (2000) Weighting and Standard Error Estimation for ABS Household Surveys, Australian Bureau of Statistics Methodology Advisory Committee Paper. Canberra.

For Benchmark 3, age strata were combined for cells where there were very few respondents. These cells were combined, within each facility, as follows:

- Across all age groups for all admitted patients
- Across all age groups in C facilities in FWLHD (Quarter 2 only)
- Across Under 18 and 18-49 age groups in C facilities in FWLHD in (Quarters 3 and 1 only)
- Across Under 18 and 18-49 age groups in C facilities in WSLHD in (Quarter 3 only)

A lower bound of one was specified in the macro. Each quarter of data was weighted separately using this process. Interim quarterly weights were used as initial response weights. Weights generated using the GREGWT macro were trimmed to 400 to avoid extreme weights. The trimming of quarterly response weights applied to 118 respondents (0.6% of all respondents).

Once four quarters of data were available, these were aggregated and the weights for facilities sampled on the basis of annual reporting were adjusted to allow reporting at the facility level. The GREGWT macro was used, in two stages, to ensure agreement of weights with populations at the margins.

Firstly, interim annual weights were calculated for the facilities sampled on the basis of annual reporting, by using the GREGWT macro with the following benchmarks.

- Benchmark 1: Facility x age stratum
- Benchmark 2: Quarter x LHD
- Benchmark 3: Quarter x age stratum
- Benchmark 4: Quarter x peer group

A lower bound of one was specified in the macro. The interim quarterly weights were used as initial response weights.

In the second stage, annual response weights were adjusted to account for disproportionate sampling of admitted emergency patients (which occurred, inadvertently, during sampling) using the GREGWT macro.

For the final annual weights, the margins were specified as follows:

- Benchmark 1: Quarter x facility (with annually-reported facilities within the same LHD combined)
- Benchmark 2: Peer group
- Benchmark 3: LHD
- Benchmark 4: Facility
- Benchmark 5: Age stratum
- Benchmark 6: Visit type
- Benchmark 7: Peer group x age stratum
- Benchmark 8: Peer group x visit type
- Benchmark 9: LHD x age stratum
- Benchmark 10: LHD x visit type
- Benchmark 11: Facility x age stratum
- Benchmark 12: Facility x visit type (with annually-reported facilities combined)

A lower bound of one was specified in the macro. Interim annual weights generated in the first stage were used as initial response weights. Weights generated using the GREGWT macro were trimmed to 400 to avoid extreme weights. The trimming of annual response weights applied to 122 respondents (0.7% of all respondents).

Analysis of weights

As part of the weighting process, an investigation of the weights is undertaken for each quarter separately to ensure that undue weight is not applied to individual responses. The two most important factors considered are the ratio of the maximum to median weight, particularly at the facility level, and the design effect.

The design effect (DEFF) was calculated for each LHD and overall, for each quarter and for the four quarters combined. The DEFF, estimated as (1+coefficient of variance (weights)²), compares the variance of estimates obtained from the stratified sample used with the variance expected for a simple random sample. Sample sizes, weighted response rates and DEFFs based on the 12 months of data are shown in Table 3 (by LHD and NSW) and Table 4 (by facility).

Table 3: Sample size, response rates and design effects (DEFF) by LHD and overall, EDPS 2014-15

LHD	Surveys Mailed	Survey Responses	Weighted Response Rate	DEFF
Central Coast	3131	755	30%	1.4
Far West	444	75	24%	1.3
Hunter New England	12605	2632	26%	1.5
Illawarra Shoalhaven	4488	1103	30%	1.5
Mid North Coast	4343	1027	29%	1.6
Murrumbidgee	3412	710	25%	1.5
Nepean Blue Mountains	2976	657	27%	1.3
Northern NSW	5301	1211	29%	1.4
Northern Sydney	6809	1927	33%	1.7
South Eastern Sydney	6294	1513	28%	1.4
South Western Sydney	7375	1442	23%	1.3
Southern NSW	2677	640	29%	1.3
St Vincent's Health Network	1640	309	25%	1.3
Sydney	4734	1134	28%	1.6
Sydney Children's Health Network	3943	1068	27%	1.3
Western NSW	5449	1104	25%	1.5
Western Sydney	5279	994	22%	1.5
NSW	80900	18301	27%	1.5

At the LHD level, the DEFFs range from just over 1.3 to 1.7. This suggests that the sample variance of estimates for some LHDs will be 1.7 times the sample variance that would have been obtained if simple random sampling had been done across the LHD. The LHDs with the largest DEFFs are those that have the greatest range in patient volumes across the facilities within the LHD. The standard errors at the LHD level are fairly small because of the sample sizes at the LHD level. Therefore the increase in standard errors caused by the survey design (and leading to a larger DEFF at LHD level) is more than offset by the fact that

each facility that is sampled has sufficient sample size to allow facility level reporting. In addition, the estimates at the LHD level have appropriate apportionment of respondents between large and small facilities. It was therefore decided not to censor larger weights further than what had already occurred by setting a global maximum weight of 400.

Table 4: Sample size, response rates and design effects (DEFF) by facility, EDPS 2014-15

Facility name	Original Peer Group	Surveys Mailed	Survey Responses	Weighted Response Rate	DEFF
Facilit	ies reported qua	arterly			
Bankstown / Lidcombe Hospital	A1	1566	292	23%	1.5
Concord Hospital	A1	1469	400	32%	1.2
Gosford Hospital	A1	1564	404	32%	1.5
John Hunter Hospital	A1	1613	392	30%	1.4
Liverpool Hospital	A1	1604	317	22%	1.2
Nepean Hospital	A1	1625	346	25%	1.3
Prince of Wales Hospital	A1	1590	322	25%	1.4
Royal North Shore Hospital	A1	1605	468	33%	1.4
Royal Prince Alfred Hospital	A1	1626	378	28%	1.5
St George Hospital	A1	1566	403	28%	1.3
St Vincent's Hospital, Darlinghurst	A1	1640	309	25%	1.3
Westmead Hospital	A1	1565	325	25%	1.2
Wollongong Hospital	A1	1568	375	29%	1.4
Sydney Children's Hospital	A2	1963	574	29%	1.2
The Children's Hospital at Westmead	A2	1980	494	25%	1.2
Calvary Mater Newcastle	A3	1533	328	27%	1.8
Sydney/Sydney Eye Hospital	A3	1583	346	27%	1.2
Auburn Hospital	В	1653	271	18%	1.3
Blacktown Hospital	В	1581	321	25%	1.5
Campbelltown Hospital	В	1639	294	22%	1.3
Canterbury Hospital	В	1639	356	24%	1.4
Coffs Harbour Base Hospital	В	1567	348	28%	1.5
Dubbo Base Hospital	В	1587	311	24%	1.3
Fairfield Hospital	В	1619	311	21%	1.3
Hornsby and Ku-Ring-Gai Hospital	В	1592	510	36%	1.5
Lismore Base Hospital	В	1539	352	28%	1.2
Maitland Hospital	В	1663	328	25%	1.3
Manly District Hospital	В	1581	381	28%	1.4
Manning Base Hospital	В	1484	394	33%	1.3
Mona Vale and District Hospital	В	1576	432	32%	1.4
Orange Health Service	В	1604	362	28%	1.2
Port Macquarie Base Hospital	В	1482	369	31%	1.5
Shoalhaven and District Memorial Hospital	В	1542	372	30%	1.3
Sutherland Hospital	В	1555	442	34%	1.4
Tamworth Base Hospital	В	1617	301	23%	1.2
The Tweed Hospital	В	1548	363	28%	1.5
Wagga Wagga Base Hospital	В	1605	330	25%	1.4
Wyong Hospital	В	1567	351	29%	1.4

Table 4: Sample size, response rates and design effects (DEFF) by facility, EDPS 2014-15 (cont.)

Facility name
Original Surveys Survey Weighted

Facilities rmidale and New England Hospital athurst Base Hospital ega District Hospital elmont Hospital owral and District Hospital	reported an C1 C1 C1	467	87	24%	
athurst Base Hospital ega District Hospital elmont Hospital	C1		87	2/10/-	4 .
ega District Hospital elmont Hospital	-	470		Z4 /0	1.4
elmont Hospital	C1	478	87	23%	1.1
•	O i	431	118	34%	1.3
owral and District Hospital	C1	441	121	35%	1.1
owrai and District Hospital	C1	443	127	36%	1.2
roken Hill Base Hospital	C1	444	75	24%	1.3
oulburn Base Hospital	C1	441	95	26%	1.2
rafton Base Hospital	C1	454	98	28%	1.1
riffith Base Hospital	C1	467	82	22%	1.2
awkesbury District Health Service	C1	465	114	32%	1.1
ount Druitt Hospital	C1	480	77	18%	1.1
urwillumbah District Hospital	C1	456	96	26%	1.1
yde Hospital	C1	455	136	35%	1.4
hellharbour Hospital	C1	456	109	29%	1.1
allina District Hospital	C2	430	105	31%	1.1
ateman's Bay District Hospital	C2	442	111	33%	1.1
ellinger River District Hospital	C2	419	108	32%	1.2
lue Mountains District Anzac Memorial Hospital	C2	443	109	31%	1.1
ulli District Hospital	C2	491	136	33%	1.1
amden Hospital	C2	504	101	24%	1.1
asino and District Memorial Hospital	C2	455	84	24%	1.2
essnock District Hospital	C2	479	76	20%	1.1
ooma Health Service	C2	441	108	30%	1.
owra District Hospital	C2	416	81	26%	1.2
eniliquin Health Service	C2	443	106	29%	1.2
orbes District Hospital	C2	456	88	23%	1.2
unnedah District Hospital	C2	456	97	26%	1.3
verell District Hospital	C2	467	90	25%	1.2
empsey Hospital	C2	456	92	25%	1.2
urri Kurri District Hospital	C2	479	85	22%	1.1
thgow Health Service	C2	443	88	26%	1.1
acksville District Hospital	C2	419	110	33%	1.2
aclean District Hospital	C2	419	113	36%	1.1
ilton and Ulladulla Hospital	C2	431	111	32%	1.1
oree District Hospital	C2	468	67	17%	1.4
oruya District Hospital	C2	430	114	32%	1.1
udgee District Hospital	C2	452	85	23%	1.2
uswellbrook District Hospital	C2	452	81	23%	1.6
arrabri District Hospital	C2	479	94	26%	1.3
•	C2		90		
arkes District Hospital		456		26%	1.2
ueanbeyan Health Service	C2	492	94	22%	1.2
ingleton District Hospital	C2	503	91	22%	1.2
umut Health Service oung Health Service	C2	441 456	102 90	29% 26%	1.3

Demographic characteristics of respondents to EDPS 2014-15

The likelihood of a patient to respond to the survey depends, at least in part, to the socio-demographic identity of the patient. For example, older patients are more likely to respond to the survey as are female patients. Furthermore, patient demographics can affect how patients respond to survey questions and the effect of differing response rates can lead to results that are not representative of the hospital's patient population. To correct for this effect, the survey program 'weights' patient responses so that the results more closely reflect the views of patients at the hospital, LHD or for NSW. The process of weighting is described in the section titled 'Weighting for data'.

Table 5 presents the demographic composition of patients by LHD, age group, visit type, peer group, Aboriginal status and gender, at each stage of the survey. Of the four columns with data:

- 1) Percentage in initial sampling frame: the percentage of patients in each category in the dataset of eligible patients, following Phase 1 screening
- 2) Percentage in sample mailed: the percentage of patients in each category provided by the NSW Ministry of Health to Ipsos for mailing, following Phase 2 screening
- 3) Percentage of respondents (unweighted): the raw/unadjusted percentage of respondents
- 4) Percentage of respondents (weighted): the weighted percentage of respondents in the final data contributing to reported results.

Table 5: Demographic characteristics of patients and EDPS respondents, 2014-15

Demographic variable	Sub-group	Percentage in patient population	% in MoH* eligible population	Percentage of respondents (Unweighted)	Percentage of respondents (Weighted)
	CCLHD	5.0	5.2	4.1	5.2
	FWLHD	0.7	0.7	0.4	0.7
	HNELHD	14.1	13.7	14.4	13.7
	ISLHD	5.9	6.0	6.0	6.0
	MLHD	3.0	3.0	3.9	3.0
	MNCLHD	4.3	4.2	5.6	4.2
	NBMLHD	4.9	5.0	3.6	4.9
	NNSWLHD	6.4	6.2	6.6	6.2
LHD	NSLHD	8.6	8.9	10.5	8.9
	SCHN	4.0	4.1	5.8	4.2
	SESLHD	8.8	9.0	8.3	9.1
	SLHD	6.4	6.2	6.2	6.2
	SNSWLHD	3.2	3.3	3.5	3.3
	SWSLHD	10.9	11.1	7.9	11.1
	SVHN	2.0	1.8	1.7	1.8
	WNSWLHD	4.8	4.6	6.0	4.6
	WSLHD	7.2	7.1	5.4	7.0
	Under 18	26.1	26.5	31.9	26.6
Age group	18-49	38.1	38.7	30.3	38.5
	50+	35.9	34.9	37.9	34.9
Visit type	Admitted Emergency	28.2	24.9	10.0	24.3
visit type	Non-admitted Emergency	71.8	75.1	90.0	75.7
	A1	34.5	35.1	25.9	35.1
	A2	4.0	4.1	5.8	4.2
D	A3	2.9	2.7	3.7	2.7
Peer group	В	33.4	33.3	41.0	33.5
	C1	12.2	12.3	7.8	12.0
	C2	13.1	12.4	15.9	12.5
Aboriginal or	No	94.0	N/A [#]	94.9	94.8
Torres Strait	Yes	5.0	N/A [#]	2.6	2.6
Islander	Unrecorded	1.0	N/A [#]	2.6	2.6
	Male	51.0	N/A [#]	48.2	47.3
Gender	Female	49.0	N/A [#]	51.8	52.7
	Unrecorded	0.0	N/A [#]	0	0

^{*}MoH = NSW Ministry of Health
#Sample summaries provided by MoH are summarised only by strata variables. As gender and Aboriginal status were not strata variables, this information was not available at this point in the process.

Reporting

Confidentiality

BHI does not receive any confidential patient information. The process of mailing of surveys and collation of responses are carried out by Ipsos Social Research Institute (Ipsos) on behalf of BHI. All personal identifiers, such as name, address etc., are removed from the data before it is provided to BHI.

Only aggregated data are published – unit record data are never published in BHI reports. To further ensure that respondents are not identifiable, BHI only publishes results that include a minimum of 30 respondents.

Statistical Analysis

Data were analysed for entire period from April 2014 to March 2015, as well as by quarter. Analyses were undertaken in SAS V9.4 using the SURVEYFREQ procedure. Strata statement variables included: facility, age and visit type.

Results were weighted for all questions except for questions related to socio-demographic characteristics and self-reported health.

For analysis of results at the quarterly level:

- Strata statement variables included: facility (with annually-reported facilities combined within LHD), LHD
 and age strata
- · Results were weighted using weights calculated for the analysis of quarterly data
- Results were generated at the NSW level, and by LHD, peer group and facility (facility-level results only reported for facilities sampled on the basis of quarterly reporting).

Where questions were comparable between years, quarterly results from the 2014-15 survey were appended to quarterly 2013-14 results. For these quarterly results, only performance-type questions are reported in *Healthcare Observer* (www.bhi.nsw.gov.au/healthcare_observer).

In Snapshot: Emergency Department Patient Report, 2014-15, statistically significant trends in the most positive category of the questions were identified using simple linear regression. A model was fitted across the eight quarters of results, weighted by the inverse of the width of the confidence interval for each point estimate. Statistically significant trends (where the p-value of the regression coefficient was less than 0.05) were only reported for questions where an LHD had a least 6 quarters of results and a coefficient of determination (R^2) of at least 0.6.

For analysis of results at the annual level:

- Strata statement variables included: facility and age strata
- Results were weighted using weights calculated for the analysis of annual data
- · Results were generated for each question in the survey

- · Results were generated at the:
 - NSW level, and by LHD, peer group and facility
 - NSW level, and by LHD, peer group and facility by triage category (Triage Categories 2,3 and Triage Categories 4,5)
 - NSW level, and by LHD, peer group and facility, by demographic characteristics outlined in Table 6
 - NSW level by triage category, by demographic characteristics outlined in Table 6.

Table 6: Demographic characteristics of EDPS respondents for reporting, 2014-15

Characteristic	Comment
Age group	0-17, 18-49, 50+ based on self-reported year of birth. Where question on year of birth was missing or invalid, administrative data were used
Gender	Male, Female. Where response were missing or invalid, administrative data were used
Education	
Main language spoken at home	Dichotomised to English, Language other than English
Rurality of hospital (NSW only)	Based on Remoteness category of postcode of location of facility
Long-standing health conditions	Dichotomised to long-standing health condition is reported and none reported for the demographic breakdown
Aboriginal status	Self-reported, dichotomised into Aboriginal or non-Aboriginal. Missing values were excluded rather than imputed from administrative source
Self-reported health status	The SF-1. Excellent, Very good, Good, Fair, Poor
Quintile of socio-economic disadvantage	Refer to the Data Dictionary: Quintile of socio-economic disadvantage
Rurality of patient residence	Based on Remoteness category of postcode of patient residence
Country of birth	Australian born vs other, derived from administrative data

Unless otherwise specified, missing responses and those who responded 'Don't know/can't remember' to questions were excluded from analysis. Typically, performance-style questions exclude missing values and 'Don't know/can't remember'-type responses. The exception is for 'Don't know/can't remember' responses for questions that ask about a third party (e.g. if family had enough opportunity to talk to doctor) or that are over 10%. Meanwhile, questions that are not related to hospital performance include results for people who responded 'Don't know/can't remember' and those who should have answered the question but did not. Results are presented only where the result was based on at least 30 respondents. For a detailed breakdown of the amount of missing or 'Don't know' responses by question, refer to Appendix 2.

Confidence intervals can be displayed in Healthcare Observer only for quarterly results. The BHI document, "Guide to Interpreting Differences" provides information in understanding comparison of results (http://www.bhi.nsw.gov.au/nsw_patient_survey_program). However, some differences in results between facilities may be due to differences in the demographic profile of patients attending those facilities. BHI is currently developing methods to standardise survey results in order to account for differences in patient mix and to optimise direct comparisons.

Calculation of percentages

The result (percentage) for each response option in the questionnaire is determined using the following method:

Numerator

The (weighted) number of survey respondents who selected a specific response option to a certain question, minus exclusions.

Denominator

The (weighted) number of survey respondents who selected any of the response options to a certain question, minus exclusions.

Calculation

= numerator/denominator X100

The results are weighted for most questions. They are not weighted for questions relating to demographics or self-reported health status.

In some cases, the results from several responses are combined to form a 'derived measure', as indicated in the reporting. For information about how these measures are developed, please see Appendix 3.

Appendix 1

Facilities included in the EDPS sampling frame

Appendix Table 1: Eligible patients, sampled patients and proportion sampled by facility, EDPS 2014-15

Facility name	Peer Group	Total eligible patients	Total sampled	Proportion sampled
Bankstown / Lidcombe Hospital	A1	37459	1566	4.2
Concord Hospital	A1	27338	1469	5.4
Gosford Hospital	A1	46315	1564	3.4
John Hunter Hospital	A1	56451	1613	2.9
Liverpool Hospital	A1	59622	1604	2.7
Nepean Hospital	A1	49162	1625	3.3
Prince of Wales Hospital	A1	38772	1590	4.1
Royal North Shore Hospital	A1	59565	1605	2.7
Royal Prince Alfred Hospital	A1	50488	1626	3.2
St George Hospital	A1	54494	1566	2.9
St Vincent's Hospital, Darlinghurst	A1	31227	1640	5.3
Westmead Hospital	A1	52020	1565	3.0
Wollongong Hospital	A1	44380	1568	3.5
Sydney Children's Hospital	A2	28869	1963	6.8
The Children's Hospital at Westmead	A2	42424	1980	4.7
Calvary Mater Newcastle	A3	22810	1533	6.7
Sydney/Sydney Eye Hospital	A3	23933	1583	6.6
Auburn Hospital	В	18103	1653	9.1
Blacktown Hospital	В	28753	1581	5.5
Campbelltown Hospital	В	46769	1639	3.5
Canterbury Hospital	В	29434	1639	5.6
Coffs Harbour Base Hospital	В	26353	1567	5.9
Dubbo Base Hospital	В	19884	1587	8.0
Fairfield Hospital	В	26092	1619	6.2
Hornsby and Ku-Ring-Gai Hospital	В	28669	1592	5.6
Lismore Base Hospital	В	21225	1539	7.3
Maitland Hospital	В	32618	1663	5.1
Manly District Hospital	В	18710	1581	8.5
Manning Base Hospital	В	18744	1484	7.9
Mona Vale and District Hospital	В	25631	1576	6.1
Orange Health Service	В	19851	1604	8.1
Port Macquarie Base Hospital	В	21147	1482	7.0
Shoalhaven and District Memorial Hospital	В	24960	1542	6.2
Sutherland Hospital	В	38566	1555	4.0
Tamworth Base Hospital	В	28407	1617	5.7
The Tweed Hospital	В	33337	1548	4.6
Wagga Wagga Base Hospital	В	25346	1605	6.3
Wyong Hospital	В	43440	1567	3.6
Armidale and New England Hospital	C1	10850	467	4.3
Bathurst Base Hospital	C1	17277	478	2.8
Bega District Hospital	C1	8615	431	5.0
Belmont Hospital	C1	18227	441	2.4
Bowral and District Hospital	C1	13139	443	3.4

Broken Hill Base Hospital	C1	12473	444	3.6
Goulburn Base Hospital	C1	11290	441	3.9
Grafton Base Hospital	C1	15189	454	3.0
Griffith Base Hospital	C1	12436	467	3.8
Hawkesbury District Health Service	C1	17159	465	2.7
Mount Druitt Hospital	C1	24149	480	2.0
Murwillumbah District Hospital	C1	10344	456	4.4
Ryde Hospital	C1	20773	455	2.2
Shellharbour Hospital	C1	20072	456	2.3
Ballina District Hospital	C2	10929	430	3.9
Bateman's Bay District Hospital	C2	9173	442	4.8
Bellinger River District Hospital	C2	3039	419	13.8
Blue Mountains District Anzac Memorial Hospital	C2	11310	443	3.9
Bulli District Hospital	C2	3661	491	13.4
Camden Hospital	C2	9531	504	5.3
Casino and District Memorial Hospital	C2	8465	455	5.4
Cessnock District Hospital	C2	11235	479	4.3
Cooma Health Service	C2	6682	441	6.6
Cowra District Hospital	C2	4177	416	10.0
Deniliquin Health Service	C2	5115	443	8.7
Forbes District Hospital	C2	4751	456	9.6
Gunnedah District Hospital	C2	5272	456	8.6
Inverell District Hospital	C2	6496	467	7.2
Kempsey Hospital	C2	13759	456	3.3
Kurri Kurri District Hospital	C2	3935	479	12.2
Lithgow Health Service	C2	7964	443	5.6
Macksville District Hospital	C2	7565	419	5.5
Maclean District Hospital	C2	7352	419	5.7
Milton and Ulladulla Hospital	C2	9813	431	4.4
Moree District Hospital	C2	4911	468	9.5
Moruya District Hospital	C2	7310	430	5.9
Mudgee District Hospital	C2	8047	452	5.6
Muswellbrook District Hospital	C2	5389	479	8.9
Narrabri District Hospital	C2	3624	456	12.6
Parkes District Hospital	C2	5440	456	8.4
Queanbeyan Health Service	C2	13329	492	3.7
Singleton District Hospital	C2	8107	503	6.2
Tumut Health Service	C2	2771	441	15.9
Young Health Service	C2	6015	456	7.6

Appendix 2

Missing and 'Don't know' responses

These data are sourced from Emergency Department Patient Survey, April 2014 to March 2015. Data are unweighted.

	Question text	Missing %	Don't know %	Missing + Don't know %*
1	What was your main form of transport to the ED?		1.3	1.3
2	Was there a problem in finding a parking place near to the ED?		0.8	0.8
3	Was the signposting directing you to the ED of the hospital easy to follow?		1.2	1.2
4	Overall, did the ambulance crew treat you with respect and dignity?	1.6	2.1	3.7
5	How would you rate how the ambulance crew and ED staff worked together?	2.4	2.6	5.0
6	Did the ambulance crew transfer information about your condition to the ED staff?	8.7	2.5	11.2
7	Overall, how would you rate the care you received from the ambulance service?	1.4	2.3	3.7
8	Were the reception staff you met on your arrival to the ED polite and courteous?	1.4	0.6	2.0
9	Did reception staff give you enough information about what to expect during your visit?	4.2	1.1	5.3
10	Did reception staff tell you how long you would have to wait for treatment?	7.8	1.8	9.6
11	Was the waiting time given to you by reception staff about right?	3.3	3.3	6.6
12	Did you experience any of the following issues when in the waiting area?		6.9	6.9
13	From the time you first arrived at the ED, how long did you wait before being triaged by a nurse - that is, before an initial assessment of your condition was made?	4.1	1.5	5.7
14	Did you stay until you received treatment?		1.6	1.6
15	Why did you leave the ED before receiving treatment?	2.1	5.0	7.1
16	After triage (initial assessment), how long did you wait before being treated by an ED doctor or nurse?	4.4	4.4	8.8
17	While you were waiting to be treated, did ED staff check on your condition?	6.5	1.3	7.8
18	While you were waiting to be treated, did your symptoms or condition get worse?	3.9	1.1	5.0
19	Did the ED doctors know your medical history, which had already been given to the triage nurse or ambulance crew?	10.8	2.6	13.4
20	Did you have confidence and trust in the ED doctors treating you?		1.0	1.0
21	Were the ED doctors polite and courteous?		1.1	1.1
22	Overall, how would you rate the ED doctors who treated you?		1.2	1.2
23	Did the ED nurses know your medical history, which had already been given to the triage nurse or ambulance crew?	10.7	3.0	13.7

24	Did you have confidence and trust in the ED nurses treating you?		0.5	0.5
25	Were the ED nurses polite and courteous?		0.7	0.7
26	Overall, how would you rate the ED nurses who treated you?		0.5	0.5
27	Did the ED health professionals introduce themselves to you?		2.3	2.3
28	Did the ED health professionals explain things in a way you could understand?		2.3	2.3
29	During your visit to the ED, how much information about your condition or treatment was given to you?		2.5	2.5
30	Were you involved, as much as you wanted to be, in decisions about your care and treatment?		2.7	2.7
31	If your family members or someone else close to you wanted to talk to the ED staff, did they get the opportunity to do so?	3.4	2.8	6.2
32	How much information about your condition or treatment was given to your family, carer or someone else close to you?	4.6	3.2	7.7
33	Were you able to get assistance or advice from an ED health professional when you needed it?		3.0	3.0
34	What was this assistance or advice required for?		7.4	7.4
35	How would you rate how the ED health professionals worked together?		3.0	3.0
36	Did you ever receive contradictory information about your condition or treatment from ED health professionals?		3.8	3.8
37	Were the ED health professionals kind and caring towards you?		2.9	2.9
38	Did you feel you were treated with respect and dignity while you were in the ED?		2.8	2.8
39	Were you given enough privacy during your visit to the ED?		3.0	3.0
40	Were your cultural or religious beliefs respected by the ED staff?		3.8	3.8
41	Did you have worries or fears about your condition or treatment while in the ED?		2.8	2.8
42	Did an ED health professional discuss your worries or fears with you?		3.2	3.2
43	Were you ever in pain while in the ED?		3.3	3.3
44	Do you think the ED health professionals did everything they could to help manage your pain?		2.6	2.6
45	Did you see ED health professionals wash their hands, or use hand gel to clean their hands, before touching you?	21.3	2.7	24.0
46	How clean were the waiting and treatment areas in the ED?		2.7	2.7
47	How safe did you feel during your visit to the ED?		2.5	2.5
48	Were there things for your child to do (such as books, games and toys)?	5.5	11.0	16.5
49	Was the area in which your child was treated suitable for someone of their age group?		10.4	10.4
50	Did the ED staff provide care and understanding appropriate to the needs of your child?		10.5	10.5
51	During your visit to the ED, did you have any tests, X-rays or scans?	2.9	5.2	8.1

52	Did an ED health professional discuss the purpose of these tests, X-rays or	1.8	1.2	3.0
53	scans with you? Did an ED health professional explain the test, X-ray or scan results in a way		1.6	1.6
54	that you could understand? What happened at the end of your visit to the ED?		3.3	3.3
55	Thinking about when you left the ED, were you given enough information		0.9	0.9
56	about how to manage your care at home? Did ED staff take your family and home situation into account when	3.8	1.5	5.3
57	planning your discharge? Thinking about when you left the ED, were adequate arrangements made by		1.3	1.3
58	the hospital for any services you needed? Did ED staff tell you who to contact if you were worried about your	11.2	1.5	12.7
59	condition or treatment after you left hospital? Thinking about your illness or treatment, did an ED health professional tell		2.0	2.0
60	you about what signs or symptoms to watch out for after you went home? Were you given or prescribed medication to take at home?		1.5	1.5
	, - ·			
61	Did an ED health professional explain the purpose of this medication in a way you could understand?		1.9	1.9
62	Did an ED health professional tell you about medication side effects to watch for?		2.6	2.6
63	Did an ED health professional tell you when you could resume your usual activities, such as when you could go back to work or drive a car?		2.1	2.1
64	Did you receive a copy of a letter from the ED doctors to your family doctor (GP)?	14.4	1.3	15.7
65	Was your departure from the ED delayed – that is, before leaving the ED to go to a ward, another hospital, home, or elsewhere?		3.8	3.8
66	Did a member of staff explain the reason for the delay? [in discharge]		4.1	4.1
67	What were the main reasons for delay? [in discharge]	4.1	4.6	8.7
68	Overall, how would you rate the care you received while in the ED?		1.3	1.3
69	If asked about your experience in the ED by friends and family how would you respond?		1.8	1.8
70	Did the care and treatment received in the ED help you?		1.6	1.6
71	In total, how long did you spend in the ED? (from when entered until left to go to a ward/another hospital/home/elsewhere)	4.5	1.9	6.4
72	Did you want to make a complaint about something that happened in the ED?		2.4	2.4
73	Why didn't you make a complaint?		2.1	2.1
74	While in the ED, did you receive, or see, any information about your rights as a patient, including how to comment or complain?	39.5	2.3	41.8
75	Not including the reason you came to the ED, during your visit, or soon afterwards, did you experience any of the following complications or problems?		3.4	3.4
76	Was the impact of this complication or problem?		2.8	2.8
77	In your opinion, were members of the hospital staff open with you about this complication or problem?		5.7	5.7
78	What year were you born?		1.7	1.7
79	What is your gender?		0.7	0.7

80	Highest level of education completed		10.4	10.4
81	Which, if any, of the following long-standing conditions do you have (including age related conditions)?		2.8	2.8
82	In general, how would you rate your health?		1.5	1.5
83	Language mainly spoken at home		1.9	1.9
84	Was an interpreter provided when you needed one in the ED?		0.8	0.8
85	Aboriginal and/or Torres Strait Islander		2.6	2.6
86	What were your reasons for going to the ED?		1.8	1.8
87	Was your visit to the ED for a condition that, at the time, you thought could have been treated by a General Practitioner (GP)?		2.2	2.2
88	In the month before visiting the ED, did you?	8.3	2.4	10.7
89	Before your visit to the ED, had you previously been to an ED about the same condition or something related to it?		2.1	2.1
90	Who completed this questionnaire?		1.4	1.4
91	Do you give permission for the Bureau of Health Information to link your answers from this survey to health records related to you (the patient)?		4.9	4.9

^{*} Percentages for this column may not equal the sum of the "Missing %" and "Don't know %" columns because they were calculated using unrounded figures.

[#] For respondents who did not answer these questions, information about age and gender were substituted with age and sex fields from administrative data (from the Health Information Exchange).

Appendix 3

Derived measures

Definition

Derived measures are those for which results are calculated indirectly from respondents' answers to a survey question. These tend to be from questions that contain a 'not applicable' type response option and are used to gather information about the array of patients' needs.

Derived measures involve the grouping together of more than one response option to a question. The derived measure 'Quintile of Disadvantage' is an exception to this rule (for more information on this, please see the appropriate Data Dictionary for this measure - http://www.bhi.nsw.gov.au/nsw_patient_survey_program).

Statistical methods

Results are expressed as the percentage of respondents who chose a specific response option or options for a question. The reported percentage is calculated as the numerator divided by the denominator (defined earlier in this technical supplement).

Results are weighted as described in this report.

Inclusions

The following questions and responses were used in the construction of the derived measures:

Derived Measure	Original Question	Derived Measure Categories	Original Question Responses		
Arrived by ambulance	What was your main	 Arrived by ambulance 	Ambulance		
	form of transport to the	Arrived other way	 Private motor vehicle 		
	ED?		Public transport		
			• Other		
Needed parking near	eded parking near Was there a problem in	Didn't need parking	Didn't need parking		
the ED	finding a parking place	Needed parking	 Yes, a big problem 		
	near to the ED?		 Yes, a small problem 		
			 No problem 		

Derived Measure	Original Question	De	rived Measure Categories	Origi	nal Question Responses
Spent time in the	Did you experience any	•	Spent time in waiting	•	Couldn't find seat
waiting area	of the following issues		area	•	Seats uncomfortable
	when in the waiting			•	Too noisy
	area?			•	Too hot
				•	Too cold
				•	Bad/unpleasant smells
				•	None of these
		•	Wasn't in waiting area	•	Wasn't in waiting area
Saw a triage nurse	From the time you first	•	Saw a triage nurse	•	Triaged immediately
•	arrived at the ED, how			•	1-15 minutes
	long did you wait before			•	16-30 minutes
	being triaged by a			•	31-59 minutes
	nurse?			•	1 hr to under 2 hrs
				•	2 hrs or more
		•	Didn't see a triage nurse	•	Didn't see a triage nurse
Received treatment	Did the ED doctors	•	Saw a doctor	•	Yes, definitely
from a doctor	know your medical		Caw a doctor	•	Yes, to some extent
	history, which had			•	No
	already been given to the triage nurse or ambulance crew?	•	Didn't see a doctor	•	Wasn't treated by a doctor
Received treatment		•	Saw a nurse	•	Yes, definitely
from a nurse	your medical history,		Saw a nuise	•	Yes, to some extent
	which had already been given to the triage nurse or ambulance crew?			•	No
		•	Didn't see a nurse	•	Wasn't treated by a nurse
Needed information	During your visit to the ED, how much	•	Needed information	•	Not enough
about condition or				•	Right amount
received treatment	information about your			•	Too much
	condition or treatment	•	Didn't need information	•	Not applicable
Wanted to be involved	was given to you? Were you involved, as much as you wanted to be, in decisions about your care and treatment?	•		•	Yes, definitely
in decisions about				•	Yes, to some extent
care and treatment				•	No
		•	Didn't want involvement	•	Not well enough to be involved
				•	No need to be involved
Had family/someone	If your family members	•	Wanted to talk to staff	•	Yes, definitely
close who wanted to			and to talk to oldli	•	Yes, to some extent
talk to staff				•	No, they did not
		•	Not applicable	•	Not applicable
Had family/someone	/someone How much information wanted about your condition or about treatment was given to	•	Wanted information	•	Not enough
close who wanted			wanted information	•	Right amount
information about				•	Too much
condition or treatment		•	Not applicable	•	It was not necessary

Derived Measure	Original Question	De	rived Measure Categories	Orig	ginal Question Responses		
Had religious or cultural beliefs to consider	Were your cultural or religious beliefs respected by the ED staff?	•	Had beliefs to consider	•	Yes, sometimes No		
Child needed things to do (such as books, games and toys)?	Were there things for your child to do (such as books, games and toys)?	•	Had beliefs to consider Child needed things to do	•	Yes, there was a lot Yes, but not enough Yes, but not for age group		
		•	Not applicable	•			
Received results of test, X-ray or scan results while in ED	Did an ED health professional explain the test, X-ray or scan	•	Received results in ED	•	Yes, to some extent No		
	results in a way that you could understand?	•	Didn't receive results	•	Not told results in ED		
Needed information on how to manage care at home	Thinking about when you left the ED, were you given enough	•	Needed information	•	Yes, to some extent		
	information about how to manage your care at home?	•	Didn't need information	•	Didn't need information		
Needed family and home situation taken into account when	Did ED staff take your family and home situation into account when planning your discharge?	•	Had situation to consider	•	Yes, to some extent		
planning discharge		•	Not necessary	•	It was not necessary		
Needed arrangements made by the hospital for services needed	Thinking about when you left the ED, were adequate arrangements	•	Needed services	•	Yes, to some extent		
after discharge	made by the hospital for any services you needed?	•	Didn't need services	•	B'II' I '		
Wanted to make a complaint about	Did you want to make a complaint about	•	Wanted to complain	•	Yes, and I did complain		
something that happened in ED	something that happened in the ED?			•	Yes, but I did not complain		
		•	Didn't want to complain	•	No		
Not including the reason came to ED, [insert complication or problem] during visit or soon afterwards	Not including the reason you came to the ED, during your visit, or soon afterwards, did you experience any of the following complications or problems?	•	 Experienced uncontrolled bleeding Experienced a negative reaction to medication Experienced complications as a result of tests or procedures Experienced a blood clot Experienced a fall 				
Highest level of education*	What is the highest level of education you have completed?	•	Less than Year 12	•	Still at school		
		•	Completed Year 12 or equivalent	•	Completed Year 12 or equivalent		
		•	Trade/tech. cert./diploma	•			
		•	University degree	•	University degree		
		•	Post grad./higher degree	•	Post grad./higher degree		

Derived Measure	Original Question	De	rived Measure Categories	Origina	al Question Responses
Has a long-standing	Which, if any, of the following long-standing conditions do you have (including age related conditions)?	•	Has condition/s	•	Full/partial deafness
condition				•	Full/partial blindness
				•	A long-standing illness
				•	A long-standing physical condition
				•	A learning disability
				•	A mental health condition
				•	A neurological condition
		•	None reported	•	None of these
				•	Missing
Needed an interpreter	Was an interpreter provided when you needed one in the ED?	•	Needed an interpreter	•	Yes, always
in the ED				•	Yes, sometimes
				•	No
		•	Didn't need interpreter	•	I didn't need an interpreter
Aboriginal and/or	Are you of Aboriginal origin, Torres Strait Islander origin or both?	•	Aboriginal	•	Aboriginal
Torres Strait Islander				•	Torres Strait Islander
origin				•	Both
		•	Non-Aboriginal	•	No
Triage Category	Triage Category	•	Triage Category 1	•	Triage Category 1
		•	Triage Categories 2,3	•	Triage Category 2
				•	Triage Category 3
		•	Triage Categories 4,5	•	Triage Category 4
				•	Triage Category 5

^{*}Respondents who were 0-16 years and reported completing a university degree or higher were excluded

Exclusions

For derived measures, the following are excluded:

- Response: 'don't know/can't remember' or similar non-committal response (with the exception of
 questions where the rate of this response was over 10% and questions that refer to the experience
 of a third party such as a family/carer)
- Response: invalid (i.e. respondent was meant to skip a question but did not)
- Response: missing (with the exception of questions that allow multiple responses or a 'none of these' option, to which the missing responses are combined to create a 'none reported' variable)

Interpretation of indicator

The higher the percentage, the more respondents fall into that response category.